NKMAXBIO We support you, we believe in your research

Human UNG antibody

Catalog Number: AUN0713

PRODUCT INFORMATION

Catalog number

AUN0713

Clone No.

k1C12

Product type

Monoclonal Antibody

UnitProt No.

P13051

NCBI Accession No.

NP 550433

Alternative Names

uracil-DNA glycosylase isoform UNG2, uracil-DNA glycosylase, UDG, UNG1, UNG2, HIGM4, uracil-DNA glycosylase 1, uracil-DNA glycosylase 2

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human UNG (1-313aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human UNG antibody

Catalog Number: AUN0713

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


The human UNG gene encodes both mitochondrial (UNG1) and nuclear (UNG2) forms of uracil-DNA glycosylase (UNG). These forms are generated from transcription from alternative promoters, promoter A and promoter B respectively, and the subsequent use of alternative splicing. UNG is responsible for the removal of uracil from DNA by hydrolysis of the N-glycosidic bond that links the base to the deoxyribose backbone, leaving an abasic site. UNG is a highly conserved enzyme found in many species.

General References

Jau der C et al.,(1991) J Bacteriol 173(1):283-290. Haug T et al.,(1998) Nucleic Acids Res. 26(6):1449-1457. Hilde N et al.,(2000) Nucleic Acids Res. 28(12):2277-2285. Stuart R.W. Bellamy et al.,(2007) Nucleic Acids Res. 35(5):1478-1487. Chih chung L et al.,(2007) J Virol 81(3):1195-1208.

DATA

Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human UNG antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: Jurkat cell lysate Lane 3.: MCF7 cell lysate Lane 4.: K562 cell lysate Lane 5.: 293T cell lysate Lane 6.: HepG2 cell lysate Lane 7.: A549 cell lysate Lane 8.: SK-OV-3 cell lysate Lane 9.: PC3 cell lysate


Immunocytochemistry/Immunofluorescence (ICC/IF)



NKMAXBio We support you, we believe in your research

Human UNG antibody

Catalog Number: AUN0713



ICC/IF analysis of UNG in HeLa cells. The cell was stained with AUN0713 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

