NKMAXBIO We support you, we believe in your research

Human 5'-Nucleotidase/CD73 antibody

Catalog Number: ATGA0554

PRODUCT INFORMATION

Catalog number

ATGA0554

Clone No.

AT89E9

Product type

Monoclonal antibody

UnitProt No.

P21589

NCBI Accession No.

NP 002517

Alternative Names

CD73, 5-NT, 5'-nucleotidase, 5'-nucleotidase ecto, Al447961, E5NT, ecto, eN, eNT, NT, NT5, Nt5e, NTE, 2210401F01Rik

Additional Information

This product was produced from tissue culture supe

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human 5'-Nucleotidase/CD73 (27-252aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, FACS

Usage

The antibody has been tested by ELISA, Western blot and FACS analysis to assure specificity and reactivity.



NKMAXBIO We support you, we believe in your research

Human 5'-Nucleotidase/CD73 antibody

Catalog Number: ATGA0554

Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


CD73 (Ecto-5-nucleotidase, 5-NT) is a membrane-bound glycoprotein that functions to hydrolyze extracellular nucleoside monophosphates into bioactive nucleoside intermediates. This enables the uptake of adenosine, inosine, and guanosine into the cell and their subsequent reconversion into ATP and GTP in the purine salvage pathway. The expression of CD73 on neuronal cells has been described during developmental processes and, on lymphocytes, CD73 serves as a maturation marker, being absent from the surface of both immature B and T cells.

General References

Sern P.C, et al., (2006) Purinergic Signal 2(2):351-360. Linda F.T, et al., (2004) J Exp Med 200(11):1395-1405. Laura A, et al., (1997) J Cell Biol 136(2):421-431.

DATA


Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human 5'-Nucleotidase/CD73 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HepG2 cell lysate Lane 2.: WiDr cell lysate Lane 3.: A431 cell lysate

Flow cytometry (FACS)



Flow cytometry analysis of 5'-Nucleotidase/CD73 in A431 cells. The cell was stained with ATGA0554 at 2-5ug for 1x10^6cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mousemonoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

