# NKMAXBIO We support you, we believe in your research

# **Human KCTD15 antibody**

Catalog Number: ATGA0178

### **PRODUCT INFORMATION**

# Catalog number

ATGA0178

#### Clone No.

AT4C3

# **Product type**

Monoclonal Antibody

#### UnitProt No.

Q96SI1

#### **NCBI Accession No.**

NP 076981

#### **Alternative Names**

BTB/POZ domain-containing protein KCTD15, MGC25497, MGC2628, potassium channel tetramerisation domain containing 15

# **PRODUCT SPECIFICATION**

#### **Antibody Host**

Mouse

# **Reacts With**

Human

#### **Concentration**

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

### **Immunogen**

Recombinant human KCTD15 (1-234) purified from E.coli

### Isotype

IgG3 kappa

#### **Purification Note**

By protein-G affinity chromatography

# **Application**

ELISA, WB, ICC/IF

#### **Usage**

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# NKMAXBio We support you, we believe in your research

# **Human KCTD15 antibody**

Catalog Number: ATGA0178

#### **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

#### **Description**


KCTD15 (Potassium channel tetramerisation domain containing 15, also known as BTB/POZ domain-containing protein KCTD15) is protein that in humans is encoded by the KCTD15 gene. KCTD15 is expressed at high level in brain and hypothalamus. The potassium channel KCTD15 was identified as a genetic loci associated with higher than normal body mass index (BMI) in humans along with genes such as GNPDA2, MTCH2, FTO, and TMEM18. Single nucleotide polymorphisms (SNPs) in non-diabetic and diabetic patients showed that FTO was most strongly associated with obesity while MTCH2 and GNPDA2 were still significantly associated with higher than normal BMI levels.

#### **General References**


Willer CJ, et al. (2010) Nat. Genet, 41(1):24-34. Elks CE, et al. (2010) PLoS Med, 7(5):e1000284. Strausberg RL, et al. (2002) Proc. Natl. Acad. Sci. U.S.A, 99(26):16899-903.

### **DATA**

### Western blot analysis (WB)



The cell lysate of HeLa (35ug) was resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human KCTD15 (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



The cell lysates (10ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human KCTD15 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1: 293T cell lysate

Lane 2: KCTD15 Transfected 293T cell lysate




# NKMAXBio We support you, we believe in your research

# **Human KCTD15 antibody**

Catalog Number: ATGA0178

# Immunocytochemistry/Immunofluorescence (ICC/IF)



Immunofluorescence of human HeLa cells stained with monoclonal anti-human KCDT15 antibody (1:500) with Texas Red (Red). Nucleus was stained by Hoechst 33342 (Blue).

