PRODUCT INFORMATION

Catalog number ATGA0206

Clone No. AT4F8

Product type Monoclonal Antibody

UnitProt No. P55263

NCBI Accession No. NP_006712

Alternative Names ADK, AK, adenosine kinase, adenosine 5'-phosphotransferase

PRODUCT SPECIFICATION

Antibody Host Mouse

Reacts With Human

Concentration 1mg/ml (determined by BCA assay)

Formulation Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen Recombinant human ADK (22-362aa) purified from E. coli

lsotype

lgG1 kappa

Purification Note By protein-G affinity chromatography

Application

ELISA,WB,ICC/IF,FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

Adenosine kinase (ADK) is the key regulator of adenosine metabolism. Because of the manifold receptordependent actions of adenosine, tight regulation of adenosine levels is crucial. The intracellular and extracellular pools of adenosine are in dynamic exchange by equilibrative and concentrative nucleoside transporters, so extracellular concentrations of adenosine are regulated by interplay of theses transporters with intracellular and extracellular enzymes of adenosine metabolism. Thus, the extracellular concentration of adenosine is enhanced by inhibition of equilibrative nucleoside transporters such as S- (4-nitrobenzyl) -6-thioinosine and cannabidiol and, stimulation of extracellular ATP breakdown and inhibition of intracellular adenosine removal.

General References

Mathews II , et al. (1998). Biochemistry 1037(45):15607-20. Lindberg B, Klenow H, Hansen K (1967). J. Biol. Chem. 242 (3): 350-6. CAPUTTO R (1951). J. Biol. Chem. 189 (2): 801-14.

DATA

Western blot analysis (WB)



The cell lysates of HepG2 and 293T (35ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human ADK (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Immunocytochemistry/Immunofluorescence (ICC/IF)



NKMAXBiO We support you, we believe in your research Human Adenosine Kinase/ADK antibody Catalog Number: ATGA0206



ICC/IF analysis of ADK in A549 cells. The cell was stained with ATGA0206 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Alexa488-anti-ADK

Flow cytometry analysis of ADK in A549 cell line, staining at 2-5ug for 1×10^{6} cells (red line). The secondary antibody used goat anti-mouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).