PRODUCT INFORMATION

Catalog number ATGA0211

Clone No. AT2E7

Product type Monoclonal Antibody

UnitProt No. P51580

NCBI Accession No. NP_000358

Alternative Names

Thiopurine S-methyltransferase, Thiopurine S-methyltransferase, TPMT, Thiopurine S-methyltransferase HGNC:12014, S adenosyl L methionine thiopurine S methyltransferase, Thiopurine methyltransferase, Thiopurine S methyltransferase.

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With Human

Concentration 1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human TPMT (1-245aa) purified from E. coli

Isotype

lgG1 kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA,WB,ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

TPMT (Thiopurine S-methyltransferase) is a very important enzyme for the metabolizm of thiopurine compounds. TPMT pharmacogenetics has been studied extensively because of its clinical significance. Many patients got benefited from the knowledge of TPMT genotype-phenotype correlation. Also, TPMT activity exhibits autosomal codominant genetic polymorphism, and patients inheriting TPMT-deficiency are at high risk of potentially fatal hematopoietic toxicity.

General References

Milstein O., et al. (2008) J Biol Chem. 283(49):34414-22. Aldape MJ., et al. (2007) J Infect Dis. 195(12):1838-45.

DATA

Western blot analysis (WB)



The cell lysate (40ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TPMT antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

The recombinant protein (50ng) and cell lysate (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with antihuman TPMT antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1. : Recombinant Protein

Lane 2. : HepG2 cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)

NKMAXBIO We support you, we believe in your research Human TPMT antibody Catalog Number: ATGA0211



ICC/IF analysis of TPMT in U87MG cells. The cell was stained with ATGA0211 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

