NKMAXBio We support you, we believe in your research

Human EGFR antibody

Catalog Number: ATGA0212

PRODUCT INFORMATION

Catalog number

ATGA0212

Clone No.

AT2H8

Product type

Monoclonal Antibody

UnitProt No.

P00533

NCBI Accession No.

NP 005219

Alternative Names

Epidermal growth factor receptor, ERBB, ERBB1, HER1, NISBD2, PIG61, mENA, ERRP, Proto-oncogene c-ErbB-1, Receptor tyrosine-protein kinase erbB-1, erythroblastic leukemia viral (v-erb-b) oncogene homolog (avian)

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human EGFR (424-605aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human EGFR antibody

Catalog Number: ATGA0212

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Epidermal growth factor receptor (EGFR) is a 170 kDa membrane protein first identified as a binding partner of EGF. Also, EGFR phosphorylated itself (autophosphorylation) in response to EGF treatment. The overexpression of EGFR in various epithelial tumors was widely reported and it has been substantiated that deregulation of EGFR itself and its signaling pathways an important role in human cancers.

General References

Cohen S, (1964) Nat Cancer Inst Monogr, 13: 13-27. Toyoda H, et al. (1995) J Biol Chem, 270: 7495-500. Carpenter G, et al. (1978) Nature, 276: 409-10.


DATA

Western blot analysis (WB)



The cell lysates of HaCaT were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human EGFR (1:250). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Immunocytochemistry/Immunofluorescence (ICC/IF)



Immunofluorescence of HaCaT cells stained with Hoechst 3342 (Blue) or phalloidin-TRITC for nucleus or actin staining, respectably. monoclonal anti-human EGFR antibody (1:250) was stained with alexa 488 (green).

