# **PRODUCT INFORMATION**

Catalog number ATGA0312

Clone No. AT4B9

**Product type** Monoclonal Antibody

**UnitProt No.** P61981

NCBI Accession No. NP\_036611

## **Alternative Names**

YWHAG, tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein gamma, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein gamma N-terminally processed, protein phosphatase 1, regulatory subunit 170, PPP1R170, 14-3- $3\gamma$ 

# **PRODUCT SPECIFICATION**

### **Antibody Host**

Mouse

**Reacts With** Human

**Concentration** 1mg/ml (determined by BCA assay)

### Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

### Immunogen

Recombinant human 14-3-3 gamma (1-247aa) purified from E. coli

### Isotype

lgG1 kappa

## **Purification Note**

By protein-A affinity chromatography

## Application

ELISA, WB, ICC/IF

### Usage

The antibody has been tested by ELISA, Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results. Recommended starting dilution is 1:1000.



#### Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

#### Description


The 14-3-3 family of proteins plays a key regulatory role in signal transduction, checkpoint control, apoptotic and nutrient-sensing pathways. 14-3-3 proteins are highly conserved and ubiquitously expressed. There are at least seven isoforms, Beta, Gamma, Epsilon, Delta, Zeta, Tau and Eta that have been identified in mammals. The 14-3-3 gamma, a subtype of the 14-3-3 family of proteins, was thought to be brain and neuron-specific. It has been shown to interact with RAF1 and protein kinase C, proteins involved in various signal transduction pathways.

### **General References**

Takagaki Y. and Manley J.L. (1992) J Biol Chem 267: 23471-23474 Takagaki Y. and Manley J.L. (2000) Mol Cell Biol 20: 1515-1525 Kleiman F.E. and Manley J.L. (1999) Science 285: 1576-1579

### DATA

#### Western blot analysis (WB)




The lysates of Hela (40ug) and Mouse brain (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with antihuman 14-3-3 gamma antibody (1:1000) Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

The recombinant proteins (50ng) were resolved by SDS-PAGE, transferred to PDVF membrane and probed with anti-human 14-3-3 gamma antibody (1:1000). Proteins were visualized using a goat antimouse secondary antibody conjugated to HRP and an ECL detection system.


- Lane 1.: Recombinant Human YWHAZ
- Lane 2.: Recombinant Human YWHAB
- Lane 3.: Recombinant Human YWHAE
- Lane 4.: Recombinant Human YWHAH
- Lane 5.: Recombinant Human YWHAG
- Lane 6.: Recombinant Human SFN
- Lane 7.: Recombinant Human YWHAQ

NKMAXBiO We support you, we believe in your research Human 14-3-3 gamma antibody Catalog Number: ATGA0312



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PDVF membrane and probed with anti-human 14-3-3 gamma antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: 293T cell lysate Lane 2.: A431 cell lysate Lane 3.: K562 cell lysate Lane 4.: NIH3T3 cell lysate

### Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of 14-3-3 gamma in HeLa cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human 14-3-3 gamma antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).