# NKMAXBIO We support you, we believe in your research

# Recombinant human IL-21 protein

Catalog Number: ATGP3802

#### PRODUCT INFORMATION

# **Expression system**

Baculovirus

#### **Domain**

30-162aa

#### UniProt No.

O9HBE4

#### **NCBI Accession No.**

NP 068575

#### **Alternative Names**

Interleukin-21 isoform 1, IL21, CVID11, IL-21, Za11

### PRODUCT SPECIFICATION

## **Molecular Weight**

16.9 kDa (145aa)

#### Concentration

0.25mg/ml (determined by Bradford assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 20% glycerol

#### **Purity**

> 85% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

#### **Biological Activity**

The activity is determined by the IFN-q ELISA in a using NK-92 cell. The ED50 range  $\leq 10 \text{ng/ml}$ .

#### Tag

His-Tag

### **Application**

SDS-PAGE, Bioactivity

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

# **BACKGROUND**

# Description

IL21, also known as interleukin-21 isoform 1, is a T-helper cytokine that regulates humoral immunity and cell-



# NKMAXBio We support you, we believe in your research

# Recombinant human IL-21 protein

Catalog Number: ATGP3802

mediated anti-tumour responses. It has broad immunoregulatory activity and can modulate both humoral and cell-mediated responses. It is important in the regulation of haematopoiesis and immune responses, and can influence lymphocyte development. It is produced by activated T cells, and it influences proliferation of T and B cells and cytolytic activity of natural killer cells. It may be a critical factor in the control of persistent viral infections. It has a potential predicating significance for survival time in patients in the development of sporadic CRC. It binds to the IL-21R and can activate Janus kinase JAK1, JAK3, STAT1, and STAT3 in EDgamma-16 cells. Recombinant human IL21, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

## **Amino acid Sequence**


<ADPEFM>QGQD RHMIRMRQLI DIVDQLKNYV NDLVPEFLPA PEDVETNCEW SAFSCFQKAQ LKSANTGNNE RIINVSIKKL KRKPPSTNAG RRQKHRLTCP SCDSYEKKPP KEFLERFKSL LQKMIHQHLS SRTHGSEDS<+H HHHHH>

#### **General References**

Parrish-Novak J., et al. (2000) Nature. 408:57-63. Cui G., et al. (2017) Clin Immunol. 183:266-272.


## **DATA**

#### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

# **Biological Activity**



The activity is determined by the IFN-g ELISA in a using NK-92 cell. The ED50 range  $\leq$  10 ng/ml.

