PRODUCT INFORMATION

Expression system HEK293

Domain 23-215aa

UniProt No. P55075

NCBI Accession No. NP_006110.1

Alternative Names

FGF8, AIGF, Androgen-induced growth factor, FGF-8, HBGF-8, Heparin-binding growth factor 8, HH6, KAL6, fibroblast growth factor 8, fibroblast growth factor 8 isoform B

PRODUCT SPECIFICATION

Molecular Weight

23.7kDa (204aa)

Concentration

0.25 mg/ml (determined by Absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol.

Purity
> 90% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells in the presence of 10 ug/ml of heparin. The ED50 range \leq 5 ug/ml.

Tag

His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.



BACKGROUND

Description

FGF-8b, also known as Fibroblast growth factor 8 isoform B, is a member of the fibroblast growth factor (FGF) family that plays a central role during prenatal development, postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. There are 4 known alternate spliced forms of FGF-8; FGF-8a, FGF-8b, FGF-8e and FGF-8f. FGF-8 is widely expressed during embryogenesis, and mediates epithelial-mesenchymal transitions. It encodes a key signaling factor, and its precise regulation is essential for embryo patterning. FGF-8b shows the strongest receptor affinity and oncogenic transforming capacity. FGF-8 signaling from the apical ectodermal ridge (AER), which borders the distal end of the limb bud, is necessary for forming normal limbs. FGF-8 is responsible for cell proliferation and survival, as well. The ability for FGF-8 to regulate cell proliferation has caused interest in its effects on tumors or squamous cell carcinoma. Recombinant human FGF-8b, fused to His-tag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

Amino acid Sequence

<DGSHM>QVTVQ SSPNFTQHVR EQSLVTDQLS RRLIRTYQLY SRTSGKHVQV LANKRINAMA EDGDPFAKLI VETDTFGSRV RVRGAETGLY ICMNKKGKLI AKSNGKGKDC VFTEIVLENN YTALQNAKYE GWYMAFTRKG RPRKGSKTRQ HQREVHFMKR LPRGHHTTEQ SLRFEFLNYP PFTRSLRGSQ RTWAPEPR<HH HHHH>

General References

Tanaka, A. et al. (1992) Proc. Natl. Acad. Sci. USA 89:8928-8932. Ornitz DM, et al. (1996). The Journal of Biological Chemistry. 271: 15292–15297. Crossley, P.H. et al. (1996) Cell, 84:127-136. Estienne A, et al.(2018) Reproduction. 155:53-62.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

Biological Activity

NKMAXBio We support you, we believe in your research Recombinant human FGF-8b protein

Catalog Number: ATGP4097

Human FGF-8b (ug/ml)

18000

16000 14000 12000 10000 RFU 8000 6000 4000 2000 0 10-2 10-1 10⁰ 10¹ 10² 10-3

Human FGF-8b stimulates cell proliferation of the Balb/3T3 mouse embryonic fibroblast cells in the presence of 10 ug/ml of heparin. The ED50 range \leq to 5 ug/ml.