# NKMAXBio We support you, we believe in your research

## Recombinant human Podocalyxin protein

Catalog Number: ATGP4090

## **PRODUCT INFORMATION**

## **Expression system**

**HEK293** 

#### **Domain**

23-429aa

#### UniProt No.

000592

#### **NCBI Accession No.**

NP 005388.2

## **Alternative Names**

PODXL, Podocalyxin, Podocalyxin isoform 2, GCTM, GCTM-2 antigen, Gp200, Podocalyxin-like protein 1, PCLP, PCLP-1, PCpodocalyxin, podocalyxin-like, PDX, gp135, PC, PODXL1

### PRODUCT SPECIFICATION

#### **Molecular Weight**

43.1kDa (416aa)

#### Concentration

1 mg/ml (determined by Absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol.

#### **Purity**

> 95% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

### Tag

His-Tag

## **Application**

SDS-PAGE

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

## **Description**

Podocalyxin, also known as PODXL, is a member of the CD34 sialomucin protein family. It is present in glomerular podocytes, endothelial cells, glandular cells in fallopian tube, uterus. A soluble form of Podocalyxin can be released into the urine of women with pre-eclampsia. It has been shown to interact with Sodium-



## NKMAXBIO We support you, we believe in your research

## Recombinant human Podocalyxin protein

Catalog Number: ATGP4090

hydrogen exchange regulatory cofactor 2. And Its interactions with L-Selectin and E-Selectin mediate the tethering of lymphocytes and metastatic tumor cells to the vascular endothelium. It is upregulated in a number of cancers and is frequently associated with poor prognosis. Based on patient survival data, high level of PODXL transcripts in tumor cells is associated with poor prognosis in renal cancer. Recombinant human Podocalyxin, fused to His-tag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

### **Amino acid Sequence**


<DGS>SPSPSPS PSQNATQTTT DSSNKTAPTP ASSVTIMATD TAQQSTVPTS KANEILASVK ATTLGVSSDS PGTTTLAQQV SGPVNTTVAR GGGSGNPTTT IESPKSTKSA DTTTVATSTA TAKPNTTSSQ NGAEDTTNSG GKSSHSVTTD LTSTKAEHLT TPHPTSPLSP RQPTSTHPVA TPTSSGHDHL MKISSSSSTV AIPGYTFTSP GMTTTLPSSV ISQRTQQTSS QMPASSTAPS SQETVQPTSP ATALRTPTLP ETMSSSPTAA STTHRYPKTP SPTVAHESNW AKCEDLETQT QSEKQLVLNL TGNTLCAGGA SDEKLISLIC RAVKATFNPA QDKCGIRLAS VPGSQTVVVK EITIHTKLPA KDVYERLKDK WDELKEAGVS DMKLGDQGPP EEAEDRFSMP < HHHHHH+>

#### **General References**

Nielsen, J.S. and K.M. McNagny (2009) J. Am. Soc. Nephrol. 20:1669-1676. Nielsen JS, McNagny KM (2008). Journal of Cell Science. 121: 3683-3692. Takeda T, et al. (2001). The Journal of Clinical Investigation. 108: 289-301.

## **DATA**

#### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

